

On Collatz theorem

Report of – September 27, 2021

Grażyna Mirkowska
Faculty of Mathematics and Natural Sciences
UKSW Wóycickiego 1/3
01-938 Warszawa Poland
G.Mirkowska@uksw.edu.pl

Andrzej Salwicki
Dombrova Research
Partyzantów 19
05-092 Łomianki, Poland
salwicki@gmail.com

Abstract. Collatz conjecture has a proof.

We present a couple of observations. 1. The problem is of algorithmic nature, The conjecture states that Collatz algorithm Cl has the halting property. 2. There is an evidence of infinite execution of the program in a non-standard model of Peano arithmetic. 3. For every natural number n there exists numbers x, y, z such that the equation $n \cdot 3^x + y = 2^z$ holds. 4. Another algorithm IC computes on triples x, y, z . The consecutive states of memory of any computation form monotone, descending sequences. 5. Hence, if a computation on triples is finite and successful then the corresponding computation of Collatz algorithm is finite too. 6. We construct an infinite set Z of elementary sentences that express the negation of halting property of Collatz algorithm. 7. The set Ax' of formulas that contains all axioms of elementary theory of addition of natural numbers and the set Z is consistent and has a model. Let \mathfrak{M} denote any structure which is a model of axioms Ax' . 8. We show that the structure \mathfrak{M} is not isomorphic to the standard structure of natural numbers with addition.

From this we infer that execution of Collatz algorithm in standard model of arithmetic is finite.

1. Introduction

The conjecture formulated by Lothar Collatz in 1937 is an algorithmic problem¹. It should be shown that the following *Cl* program, executed in the standard structure \mathfrak{N} of natural numbers with addition² has a finite computation for each $n \neq 0$.

We will be investigating the stop property of the following *Cl* program.

$$Cl : \left\{ \begin{array}{l} \text{while } n \neq 1 \text{ do} \\ \quad \text{if } even(n) \text{ then } n := \frac{n}{2} \text{ else } n := 3n + 1 \text{ fi} \\ \text{od} \end{array} \right\}$$

In 2004, we noticed that there is a counterexample, see Appendix C (page 29). The two conclusions that can be made out of it, are:

- The formulation of the Collatz hypothesis requires clarification that the calculations are carried out in the standard model of natural numbers with addition. Note that, the operations of multiplication by 3 and division by 2 can be defined in Presburger arithmetic by means of addition.
- The Peano axioms, much less the Presburger axioms, are not sufficient to prove the conjecture.

And we found that if the conjecture is true, then it is a theorem of the algorithmic theory of natural numbers $AT\mathcal{N}$, (see the page 4).

1.1. Highpoints of the proof

They can be listed as follows

- the problem is of algorithmic nature (halting property).
- It is easy to construct an algorithmic formula that expresses the halting property of Collatz program. It is difficult to prove the formula.
- The complement of Collatz tree, if it is a non-empty set, is a rich, tree-like structure of infinitely many elements.

¹It may be surprising that some people place this problem in the theory of dynamical systems.

²programmers may prefer another formulation: *execution of CL algorithm in unsigned integers of unlimited precision*

- Every natural number can be represented using three natural numbers, $\forall_n, \exists_{x,y,z} n \cdot 3^x + y = 2^z$.
- three easy observations
 1. For a given number n , if the computation of Collatz algorithm is finite, then there exists triple $\langle x, y, z \rangle$ such that the triple represents n and the execution of algorithm IC is finite.
 2. if there is a triple $\langle x, y, z \rangle$ such that the equality $\forall_n, \exists_{x,y,z} n \cdot 3^x + y = 2^z$ holds and the execution of program IC is finite, then the execution of Collatz algorithm is finite.
 3. if the execution of program IC is infinite and the equality $\forall_n, \exists_{x,y,z} n \cdot 3^x + y = 2^z$ holds, then the execution of Collatz algorithm for number n is infinite
- If for an element n the execution of Collatz algorithm is infinite then there exists a triple $\langle x, y, z \rangle$ such that the equality $\forall_n, \exists_{x,y,z} n \cdot 3^x + y = 2^z$ holds and the execution of program IC is infinite

2. Halting formula

Halting formula of a program K is any formula χ that expresses the finiteness of a computation of the program K .

Note, for many programs, their halting formulas are beyond the language of first-order logic. However, for every program its halting property can be expressed by an algorithmic formula of program calculus (i.e. algorithmic logic).

The following formula (halt) expresses the halting property of Collatz program Cl . It is to be shown that the formula is valid in the structure \mathfrak{N} of natural numbers with addition.

$$\left\{ \begin{array}{l} \text{while } n \neq 1 \text{ do} \\ \quad K : \text{if } even(n) \text{ then } n := \frac{n}{2} \text{ else } n := 3n + 1 \text{ fi} \\ \quad \text{od} \end{array} \right\} (n = 1) \quad (\text{halt})$$

Formula (LC) that asserts, there exists an iteration of the program K , such that the condition $(n = 1)$ holds after execution of program

K^i is a halting formula of program Cl too.

$$\cup \left\{ \begin{array}{l} \text{if } n \neq 1 \text{ then} \\ \quad \left[\begin{array}{l} \text{if } n \bmod 2 = 1 \\ \quad \text{then } n \leftarrow 3n + 1 \\ \quad \text{else } n \leftarrow n \bmod 2 \end{array} \right] \\ \text{fi} \\ \text{fi} \end{array} \right\} (n = 1) \quad (\text{LC})$$

Our goal is to show that the halting formula is valid in the standard structure \mathfrak{N} of natural numbers or to prove the formula in algorithmic theory of natural numbers \mathcal{ATN} . For any model of the theory is isomorphic to the structure \mathfrak{N} . See [MS87], page 139.

$$\text{Axioms of } \mathcal{ATN} \left\{ \begin{array}{l} \forall_x x + 1 \neq 0 \\ \forall_{x,y} x + 1 = y + 1 \implies x = y \\ \forall_x \{y := 0; \text{while } y \neq x \text{ do } y := y + 1 \text{ od}\} (y = x) \end{array} \right\} \quad (\text{ATN})$$

Theory \mathcal{ATN} can be extended, we can add definitions of useful operations $+, \times 3, \div 2$, and the parity unary relation. We shall use the following notation the atomic formula $e(n)$ reads n is even, and formula $o(n)$ reads n is odd.

Making use of axioms of calculus of programs \mathcal{AL} and axioms of algorithmic theory of natural numbers \mathcal{ATN} we can write a couple of formulas that are equivalent to the halting formula halt .

Below we display one of these formulas

$$\left\{ \begin{array}{l} (o(n) \wedge \mathbf{n} = 1) \vee \\ (e(n) \wedge o(\frac{n}{2}) \wedge \mathbf{n} = 2) \vee \\ (e(n) \wedge e(\frac{n}{2}) \wedge o(\frac{n}{4}) \wedge \mathbf{n} = 4) \vee \\ (e(n) \wedge e(\frac{n}{2}) \wedge e(\frac{n}{4}) \wedge o(\frac{n}{8}) \wedge \mathbf{n} = 8) \vee \\ (e(n) \wedge e(\frac{n}{2}) \wedge e(\frac{n}{4}) \wedge e(\frac{n}{8}) \wedge o(\frac{n}{16}) \wedge \mathbf{n} = 16) \vee \\ \left(\begin{array}{l} e(n) \wedge e(\frac{n}{2}) \wedge e(\frac{n}{4}) \wedge e(\frac{n}{8}) \wedge e(\frac{n}{16}) \wedge o(\frac{n}{32}) \wedge \mathbf{n} = 32) \vee \\ o(n) \wedge e(3n+1) \wedge e(\frac{3n+1}{2}) \wedge e(\frac{3n+1}{4}) \wedge e(\frac{3n+1}{8}) \wedge o(\frac{3n+1}{16}) \wedge \mathbf{n} = 5 \end{array} \right) \vee \\ \left(\begin{array}{l} (e(n) \wedge e(\frac{n}{2}) \wedge e(\frac{n}{4}) \wedge e(\frac{n}{8}) \wedge e(\frac{n}{16}) \wedge e(\frac{n}{32}) \wedge o(\frac{n}{64}) \wedge \mathbf{n} = 64) \vee \\ e(n) \wedge o(\frac{n}{2}) \wedge e(\frac{3n+1}{2}) \cdots \wedge \mathbf{n} = 10 \end{array} \right) \vee \\ \left(\begin{array}{l} e(n) \wedge e(\frac{n}{2}) \wedge e(\frac{n}{4}) \wedge e(\frac{n}{8}) \wedge e(\frac{n}{16}) \wedge e(\frac{n}{32}) \wedge e(\frac{n}{64}) \wedge o(\frac{n}{128}) \wedge \mathbf{n} = 128) \vee \\ o(n) \wedge e(3n+1) \wedge e(\frac{3n+1}{2}) \cdots \wedge \mathbf{n} = 21 \end{array} \right) \vee \\ \left(\begin{array}{l} e(n) \wedge o(\frac{n}{2}) \wedge e(\frac{3n+1}{2}) \cdots \wedge \mathbf{n} = 20 \vee \\ o(n) \wedge e(3n+1) \wedge o(\frac{3n+1}{2}) \cdots \wedge \mathbf{n} = 3 \end{array} \right) \vee \\ \left\{ \begin{array}{l} \text{if } n \neq 1 \text{ then} \\ \quad \left[\begin{array}{l} \text{if } n \bmod 2 = 1 \\ \quad \text{then } n \leftarrow 3n + 1 \\ \quad \text{else } n \leftarrow n \bmod 2 \\ \quad \text{fi} \end{array} \right] \\ \text{fi} \end{array} \right\}^8 \cup \left\{ \begin{array}{l} \text{if } n \neq 1 \text{ then} \\ \quad \left[\begin{array}{l} \text{if } n \bmod 2 = 1 \\ \quad \text{then } n \leftarrow 3n + 1 \\ \quad \text{else } n \leftarrow n \bmod 2 \\ \quad \text{fi} \end{array} \right] \\ \text{fi} \end{array} \right\} (n = 1) \end{array} \right\}$$

Based on these experiments, we can make some observations:

- There are many equivalent formulas that express the halting property of Collatz algorithm. In particular, each formula δ_i from the sequence $\{\delta_i\}$, $i = 0.1.2.\dots$

$$\bigvee_{j=0}^i \{K^j\}(n = 1) \vee \{K^{i+1}\} \bigcup \{K\}(n = 1) \quad \text{where } i = 0.1.2. \quad (\delta_i)$$

is a halting formula of Collatz algorithm.

Evidently, the property can be expressed by other formulas.

- There is an easy way to create the i -th formula of this sequence

$$\delta_0 \stackrel{df}{=} (o(n) \wedge \mathbf{n} = 1)$$

$$\delta_{i+1} \stackrel{df}{=} \{K\}\delta_i$$

Now, by distributivity of algorithms over disjunction (i.e. axiom Ax₁₅) one can obtain

$$\{K\}\delta_i \equiv \{K\}\left(\bigvee_{j=0}^i \{K^j\}(n=1) \vee \{K^{i+1}\} \bigcup \{K\}(n=1)\right) \quad (1)$$

$$\equiv \left(\bigvee_{j=0}^{i+1} \{K^j\}(n=1) \vee \{K^{i+2}\} \bigcup \{K\}(n=1)\right) \quad (2)$$

$$\equiv \delta_{i+1} \quad (3)$$

- Let an algebraic structure \mathfrak{A} has the signature consistent with the signature of elementary theory of natural numbers with addition, let v be a valuation of variable n . From the definition of semantics we have

$$(\delta_i)_{\mathfrak{A}}(v) = l.u.b. \left(\bigvee_{j=0}^i \{K^j\}(n=1) \right)_{\mathfrak{A}}(v)$$

- the correspondence between the levels of the Collatz tree and the components of the halting formula is clearly visible,
- at higher levels of the Collatz tree more and more odd numbers appear - and the subsequent components of the halting formula are alternatives of longer and longer conjunctions of factors $o(\cdot)$ or $e(\cdot)$,
- Note, the alternative $(n=128 \vee n=21 \vee n=20 \vee n=3)$ may be written as follows $(n \cdot 3^0 + 0 = 2^7 \vee n \cdot 3^1 + 1 = 2^6 \vee n \cdot 3^1 + 4 = 2^6 \vee n \cdot 3^2 + 5 = 2^5)$.

In the section 4 we shall use this observation.

3. Collatz tree

It is easy to notice that the set of those natural numbers for which the computation of the Collatz algorithm is finite, forms a tree.

Definition 3.1. *Collatz tree* \mathcal{DC} is a subset $D \subset N$ of the set N of natural numbers and the function f defined on the set $D \setminus \{0, 1\}$.

$$\mathcal{DC} = \langle D, f \rangle$$

where $D \subset N, 1 \in D, f: D \setminus \{0, 1\} \rightarrow D$.

Function f is determined as follows

$$f(n) = \begin{cases} n \div 2 & \text{when } n \bmod 2 = 0 \\ 3n + 1 & \text{when } n \bmod 2 = 1 \end{cases}$$

, the set D is the least set containing the number 1 and closed with respect to the function f ,

$$D = \{n \in N : \exists_{i \in N} f^i(n) = 1\}.$$

As it is easy to see, this definition is highly entangled and the decision whether the set D contains every natural number is equivalent to the Collatz problem.

Remark 3.1. Set D has the following properties :

$$x \in D \implies (x + x) \in D \quad (4)$$

$$x \in D \wedge \exists_y x = y + y \implies y \in D \quad (5)$$

$$x \in D \wedge \exists_y x = y + y + 1 \implies (x + x + x + 1) \in D \quad (6)$$

$$x \in D \wedge (\exists_e \exists_z e = z + z + 1 \wedge x = e + e + e + 1) \implies e \in D \quad (7)$$

Implications (4) and (7) show left and right son of element x .

Similar, interesting properties has the complement of set D , if it is a non-empty set. Let $cD \stackrel{df}{=} N \setminus D$ denote the complement of set D .

Remark 3.2. If the complement $N \setminus D$ is a non-empty set, then it has similar properties:

$$x \in cD \implies (x + x) \in cD \quad (8)$$

$$x \in cD \wedge \exists_y x = y + y \implies y \in cD \quad (9)$$

$$x \in cD \wedge \exists_y x = y + y + 1 \implies (x + x + x + 1) \in cD \quad (10)$$

$$x \in cD \wedge (\exists_e \exists_z e = z + z + 1 \wedge x = e + e + e + 1) \implies e \in cD \quad (11)$$

Note, both sets D and cD may be considered as graphs. Their structures are similar. However, the graph cD is not a tree .

Remark 3.3. If Collatz conjecture is not true, then both sets D and $N \setminus D$ are infinite.

From properties (6) and (7) follows the

Fact 3.1. If an x element does not belong to the Collatz tree then the computation of the Collatz algorithm starting with the state $v(n) = x$ is not finite.

Let us terminate this section with the following observation

Remark 3.4. In a non-standard model of elementary theory of natural numbers with addition, the complement of Collatz tree is an infinite set.

For in this model there are unreachable elements, c.f. section 9.

4. Triples

The observations made when analyzing the halting formula of Collatz algorithm inspired us to introduce the notion of triple representing a given natural number n . We shall also discuss the computations "on triples". Let us begin with the following remark

Fact 4.1. *For every natural number $n \neq 0$ there exist many triples x, y, z of natural numbers such that the following equality holds*

$$n \cdot 3^x + y = 2^z$$

We say that triple x, y, z represents the number n and denote it by $\langle x, y, z \rangle \asymp n$.

Proof:

The proof of this intuitive fact uses the law of Archimedes ³.

Let n be an arbitrary natural number. We choose a number x , it may be an arbitrary natural number. Let the number $k = n \cdot 3^x$. Put $z = (\mu z)(2^z \geq k)$. and $y = 2^z - n \cdot 3^x$. Obviously the equality $n \cdot 3^x + y = 2^z$ holds. \square

Example 4.1. Triple $\langle 1, 7, 6 \rangle$ represents number 19 for $19 \cdot 3 + 7 = 2^6$. Number 19 is represented by many more triples.

³We recall that the law is not an elementary property, it can be formulated by an algorithmic formula $\forall_{0 < a < b} \{ z := a; \text{ while } z \leq b \text{ do } z := z + a \text{ od} \} (z > b) \}$.

$\langle 1, 7, 6 \rangle$	$19 \cdot 3^1 + 7 = 2^6$
$\langle 2, 85, 8 \rangle$	$19 \cdot 3^2 + 85 = 2^8$
$\langle 3, 511, 10 \rangle$	$19 \cdot 3^3 + 511 = 2^{10}$
$\langle 4, 509, 11 \rangle$	$19 \cdot 3^4 + 509 = 2^{11}$
$\langle 5, 3575, 13 \rangle$	$19 \cdot 3^5 + 3575 = 2^{13}$
$\langle 6, 2533, 14 \rangle$	$19 \cdot 3^6 + 2533 = 2^{14}$
$\langle 7, 23983, 16 \rangle$	$19 \cdot 3^7 + 23983 = 2^{16}$
...	

Note, not every triple represents a number, consider $\langle 2, 4, 11 \rangle$, $\langle 2, 4044, 11 \rangle$.

Fact 4.1 is a theorem of elementary theory of natural numbers with addition \mathcal{T} , c.f. section 8.

Theorem 4.1. The sentence $\forall_n \exists_{x,y,z} n \cdot 3^x + y = 2^z$ is a theorem of theory \mathcal{T} ,

The proof is in section 8. Hence the theorem is valid in any model of theory \mathcal{T} ,

4.1. Properties of triples

Definition 4.1. Triples $\langle x, y, z \rangle$ and $\langle u, v, t \rangle$ are equivalent if they represent the same natural number .

$$\langle x, y, z \rangle \equiv \langle u, v, t \rangle \quad \stackrel{\text{df}}{=} \quad \frac{2^z - y}{3^x} \in N \wedge \frac{2^z - y}{3^x} = \frac{2^t - v}{3^u}$$

Moreover, one can define a (lexicographical) order in the set of triples \succ .

Definition 4.2.

$$\langle u, v, t \rangle \succ \langle x, y, z \rangle \stackrel{\text{df}}{\Leftrightarrow} x < u \text{ lub } x = u \text{ i } z < t \text{ lub } x = u \text{ i } z = t \text{ i } y < v$$

It follows from the fact 4.1 that , for every natural number n there is a triple representing n such, that the number y is bigger than an arbitrarily chosen number k .

Definition 4.3. *The relation of parity of a triple is determined by the parity of number y .*

$$\text{odd}(\langle x, y, z \rangle) \stackrel{\text{df}}{\Leftrightarrow} y \text{ jest nieparzyste}$$

Another fact

Fact 4.2. *Number n is odd iff the triple $\langle x, y, z \rangle$ that represents n is odd .*

Definition 4.4. One

$$\text{equal1}(\langle x, y, z \rangle) \stackrel{\text{df}}{\Leftrightarrow} (2^z - y = 3^x)$$

Definition 4.5. Klasa trójków równowaznych jest zbiorem trójków reprezentujących tę samą liczbę n .

Jedynka jest reprezentowana przez wiele trójków, np. $\langle 0, 0, 0 \rangle, \langle 0, 1, 1 \rangle, \langle 1, 1, 2 \rangle, \langle 1, 5, 3 \rangle, \langle 2, 1, 4 \rangle$

Na trójkach możemy określić dwie operacje

Definition 4.6. Operacja div2 przeprowadza trójkę parzystą $\langle x, y, z \rangle$ w trójkę $\langle x, y \div 2, z - 1 \rangle$

$$\langle x, y, z \rangle \xrightarrow{\{\text{div2}\}} \langle x, y \div 2, z - 1 \rangle$$

Definition 4.7. Operacja mult3 określona jest na trójkach $\langle x, y, z \rangle$ takich, że trójka $\langle x, y, z \rangle$ jest nieparzysta i $x > 0$ i $y > 3^{x-1}$. W takim przypadku operacja mult3 przeprowadza trójkę $\langle x, y, z \rangle$ w trójkę $\langle x - 1, y - 3^{x-1}, z \rangle$

$$\langle x, y, z \rangle \xrightarrow{\{\text{mult3}\}} \langle x - 1, y - 3^{x-1}, z \rangle$$

Zauważ

Fact 4.3. Trójka $\langle x, y, z \rangle$ reprezentuje liczbę parzystą n wttw gdy trójka $\langle x, y \div 2, z - 1 \rangle$ reprezentuje liczbę $n \div 2$.

Ponadto $\langle x, y \div 2, z - 1 \rangle \prec \langle x, y, z \rangle$.

Jeśli liczba y jest nieparzysta i zachodzi $x > 0 \wedge y > 3^{x-1}$ to trójka $\langle x, y, z \rangle$ reprezentuje liczbę n wttw gdy trójka $\langle x - 1, y - 3^{x-1}, z \rangle$ reprezentuje liczbę $3n + 1$.

Ponadto $\langle x - 1, y - 3^{x-1}, z \rangle \prec \langle x, y, z \rangle$.

Zauważmy z kolej

Lemma 4.1. *Jeśli program K : {if odd then mult3 else div2 fi} przeprowadza trójkę $\langle x, y, z \rangle$ w trójkę $\langle u, v, t \rangle$, to wynikowa trójka $\langle u, v, t \rangle$ jest mniejsza od trójki $\langle x, y, z \rangle$.*

$$\langle x, y, z \rangle \xrightarrow{\text{if odd then mult3 else div2 fi}} \langle u, v, t \rangle \text{ implikuje } \langle u, v, t \rangle \prec \langle x, y, z \rangle$$

Proof:

Jeśli liczba y jest parzysta to od z odejmujemy jedynkę i dzielimy y przez 2. W przeciwnym przypadku zmniejszana jest liczba x i od y odejmowana jest liczba 3^x . \square

W każdym kroku obliczenia algorytmu Collatza wyrażenie $x + z$ zmniejsza swoją wartość o 1, zmniejszana jest też wartość zmiennej y .

Wynika stąd, że każde obliczenie w strukturze \mathfrak{T} jest skończone. Obserwacje poczynione dotychczas pozwalają stwierdzić, że następujący diagram jest przemienny.

Lemma 4.2. *Działanie na trójkach prowadzi od trójki $\langle x, y, z \rangle$ kodującą liczbę n i spełnajacej przy tym warunek $\zeta : ((y \bmod 2 = 1) \Rightarrow (x > 0 \wedge y > 3^{x-1}))$, do kolejnej trójki $\langle u, v, t \rangle$, która reprezentuje wynik m następującej instrukcji warunkowej $K : \{\text{if odd}(n) \text{ then } n := 3 * n + 1 \text{ else } n := n \text{ div } 2 \text{ fi}\}$.*




Diagram ten jest przemienny pod warunkiem, że można wykonać operację odejmowania tzn. gdy y jest liczbą parzystą lub y jest nieparzyste i ($x > 0$ lub $y > 3^{x-1}$). Semantyczną treść opisaną powyższym diagramem można wyrazić następującą formułą

$$((n \cdot 3^x + y = 2^z \wedge \text{odd}(y)) \Rightarrow (x > 0 \wedge y > 3^{x-1})) \Rightarrow \{K; \bar{K}\}(n \cdot 3^x + y = 2^z)$$

Pamiętajmy, że programy K oraz \bar{K} są niezależne, a więc przemiennne.

$$((n \cdot 3^x + y = 2^z \wedge \text{odd}(y)) \Rightarrow (x > 0 \wedge y > 3^{x-1})) \Rightarrow \{\bar{K}; K\}(n \cdot 3^x + y = 2^z)$$

Przemiennosć tego diagramu pozwala nam dostrzec, że każdemu obliczeniu algorytmu Collatza Cl w strukturze \mathfrak{N} liczb naturalnych, odpowiada obliczenie sprzężonego z Cl , algorytmu IC w strukturze trójek \mathfrak{T} .

Lemma 4.3. If the following condition is satisfied θ

$$\theta : (n \cdot 3^x + y = 2^z) \wedge \bigcap \left\{ \begin{array}{l} \text{if } 3^x + y \neq 2^z \\ \text{then} \\ \quad \text{if } odd(y) \\ \quad \text{then } x, y := x - 1, y - 3^{x-1} \\ \quad \text{else } z, y := z - 1, y \div 2 \\ \text{fi} \\ \text{fi} \end{array} \right\} (odd(y) \implies (x > 0 \wedge y > 3^{x-1}))$$

Then the following diagram commutes

$$\begin{array}{ccc} n & \xrightarrow{\{\text{while } n \neq 1 \text{ do if } odd(n) \text{ then } n := 3n+1 \text{ else } n := n/2 \text{ fi od}\}_{\mathfrak{N}}} & 1 \\ \uparrow n \cdot 3^x + y = 2^z & & \uparrow 3^x + y = 2^z \\ \langle x, y, z \rangle & \xrightarrow{\left\{ \begin{array}{l} \text{while } 3^x + y \neq 2^z \text{ do} \\ \quad \text{if } odd(y) \text{ then } x, y, z := x - 1, y - 3^{x-1}, z \\ \quad \text{else } x, y, z := x, y/2, z - 1 \text{ fi} \\ \text{od} \end{array} \right\}_{\mathfrak{T}}} & \langle x, y, z \rangle \end{array} \quad (\text{CCI})$$

This property is expreses by the formula

$$\theta \Rightarrow (\{Cl\}(n = 1) \Leftrightarrow \{IC\}(3^x + y = 2^z)) . \quad (\text{RCI})$$

Proof:

goes by induction with respect to the number of iterations. Base of induction reduces to the commutativity of the preceding fact

$$\zeta \Rightarrow \left[\begin{array}{ccc} n & \xrightarrow{\{K\}_{\mathfrak{N}}} & m \\ n \cdot 3^x + y = 2^z \uparrow & & \uparrow m \cdot 3^u + v = 2^t \\ \langle x, y, z \rangle & \xrightarrow{\bar{K}_{\mathfrak{T}}} & \langle u, v, t \rangle \end{array} \right] .$$

If the execution of algorithm is longer, say of length 2, then the following diagram applies

$$(\zeta \wedge \{\bar{K}\}\zeta) \Rightarrow \left[\begin{array}{ccccc} n & \xrightarrow{\{K\}_{\mathfrak{N}}} & n_1 & \xrightarrow{\{K\}_{\mathfrak{N}}} & n_2 \\ n \cdot 3^x + y = 2^z \uparrow & & n_1 \cdot 3^{x_1} + y_1 = 2^{z_1} \uparrow & & n_2 \cdot 3^{x_2} + y_2 = 2^{z_2} \uparrow \\ \langle x, y, z \rangle & \xrightarrow{\bar{K}_{\mathfrak{T}}} & \langle x_1, y_1, z_1 \rangle & \xrightarrow{\bar{K}_{\mathfrak{T}}} & \langle x_2, y_2, z_2 \rangle \end{array} \right] .$$

By induction, we prove that for every natural number i the following diagram commutes

$$\bigwedge_{j=0}^i \{\bar{K}\}^j \zeta \Rightarrow \left| \begin{array}{ccccccccc} n & \xrightarrow{\{K\}_{\mathfrak{N}}} & n_1 & \xrightarrow{\{K\}_{\mathfrak{N}}} & n_2 & \xrightarrow{\dots} & n_{i-1} & \xrightarrow{\{K\}_{\mathfrak{N}}} & n_i \\ n \cdot 3^x \uparrow + y = 2^z & & n_1 \cdot 3^{x_1} + y_1 \uparrow = 2^{z_1} & & n_2 \cdot 3^{x_2} + y_2 \uparrow = 2^{z_2} & & \dots \uparrow & & n_i \cdot 3^{x_i} + y_i = 2^{z_i} \uparrow \\ \langle x, y, z \rangle & \xrightarrow{\bar{K}_{\mathfrak{T}}} & \langle x_1, y_1, z_1 \rangle & \xrightarrow{\bar{K}_{\mathfrak{T}}} & \langle x_2, y_2, z_2 \rangle & \xrightarrow{\dots} & \langle x_{i-1}, y_{i-1}, z_{i-1} \rangle & \xrightarrow{\bar{K}_{\mathfrak{T}}} & \langle x_i, y_i, z_i \rangle \end{array} \right|$$

Now, if for some $i \in N$ the number $n_i = 1$, then $\forall_k n_{i+k} = 1$. Thus, the formula RCI has a proof. \square

Trzy łatwe i jeden nieoczywisty lemat.

Lemat 1. Jeśli dla pewnego elementu n obliczenie algorytmu Collatza jest skończone to istnieją trzy elementy x, y, z takie, że zachodzi równość $n \cdot 3^x + y = 2^z$ i obliczenie algorytmu IC dla tej trójki jest skończone i wolne od błędu.

Lemat 2. Jeśli istnieje trójką x, y, z reprezentująca element n i obliczenie algorytmu trójkowego jest skończone i wolne od błędu Err , to obliczenie algorytmu Collatza dla n jest skończone.

Lemat 3. W każdej strukturze algebraicznej \mathfrak{A} będącej modelem aksjomatów elementarnej teorii dodawania liczb naturalnych, jeśli istnieje trójką x, y, z reprezentująca element n i obliczenie algorytmu trójkowego jest nieskończone, to obliczenie algorytmu Collatza dla n jest nieskończone.

Lemat 4. W każdej strukturze algebraicznej \mathfrak{A} będącej modelem aksjomatów elementarnej teorii dodawania liczb naturalnych, jeśli dla pewnego elementu n obliczenie algorytmu Collatza jest nieskończone to istnieją trzy elementy x, y, z takie, że zachodzi równość $n \cdot 3^x + y = 2^z$ i obliczenie algorytmu IC dla tej trójki jest nieskończone i wolne od błędu.

The same conclusion may be obtained by an almost formal proof of the formula

$$\theta \wedge \neg\{n := \varepsilon; Cl\}(n = 1) \Rightarrow \neg\{IC\}true$$

Myszę, że da się dowieść takiego zdania (z meta-teorii \mathcal{ATN} dla każdej liczby naturalnej $i = o, 1, \dots$ twierdzeniem teorii \mathcal{ATN} jest zdanie o postaci

$$\varepsilon 3^x + y = 2^z \wedge y_i \wedge \{n := \varepsilon; K^i\}(n \neq 1) \Rightarrow \{\bar{K}^i\}(3^x + y \neq 2^z)$$

Jeśli tak jest to stosując dwukrotnie aksjomat Ax_{17} otrzymamy stwierdzenie, że twierdzeniem teorii \mathcal{ATN} jest każda formuła $(n3^x + y = 2^z \wedge \bigcap\{K\}(oddy \rightarrow) \wedge \neg\{Cl\}(n = 1) \Rightarrow \{\bar{K}\}^i(3^x + y \neq 2^z)$, gdzie $i = 0, 1, \dots$. Stąd przez zastosowanie reguły wnioskowania (r5) otrzymamy

$$n3^x + y = 2^z \wedge \bigcap\{K\}(oddy) \wedge \neg\{Cl\}(n = 1) \Rightarrow \neg\{IC\}(Err \vee 3^x + y = 2^z)$$

4.2. Zbiory T_n

Każdej liczbie n przyporządkowujemy zbiór T_n trójkę x, y, z takich, że zachodzi równość $n3^x + y = 2^z$.

Zbiór taki jest niepusty.

Zbiory T_n są parami rozłączne, tj. $n \neq m \implies T_n \cap T_m = \emptyset$.

Zbiór T_n jest zamknięty ze względu na operacje $o_1, o_2, o_3, o_4, o_5, o_6$

$$o_1(\langle x, y, z \rangle) = \langle x + 1, 3y + 2^z, z + 2 \rangle \tag{12}$$

$$o_2(\langle x, y, z \rangle) = \langle x + 1, 3y - 2^z, z + 1 \rangle \quad \text{gdy } 3y > 2^z \tag{13}$$

$$o_3(\langle x, y, z \rangle) = \langle x, y + 2^z, z + 1 \rangle \tag{14}$$

$$o_4(\langle x, y, z \rangle) = \langle x, y - 2^{z-1}, z - 1 \rangle \quad \text{gdy } y > 2^{z-1} \tag{15}$$

$$o_5(\langle x, y, z \rangle) = \langle x - 1, \frac{y - 2^{z-2}}{3}, z - 2 \rangle \quad \text{gdy } y > 2^{z-2} \text{ i } (y - 2^{z-2}) \bmod 3 = 0 \tag{16}$$

$$o_6(\langle x, y, z \rangle) = \langle x - 1, \frac{y + 2^{z-1}}{3}, z - 1 \rangle \tag{17}$$

Wynika stąd, że zbiór T_n jest nieskończony i nieograniczony.

Zauważ związki

$$o_5(o_1(\langle x, y, z \rangle)) = \langle x, y, z \rangle$$

$$o_4(o_3(\langle x, y, z \rangle)) = \langle x, y, z \rangle$$

$$o_6(o_2(\langle x, y, z \rangle)) = \langle x, y, z \rangle$$

$$o_4(o_1(\langle x, y, z \rangle)) = \langle x, y, z \rangle$$

O trójkach zbędnych (niepotrzebnych) ...

Trójka większa od trójki przydatnej nie musi być przydatna, może być zbędna.

PRZYKŁAD.

Ale też od każdej trójki dobrej istnieje większa od niej trójka dobra.

4.3. Program IC

A więc można realizować obliczenia algorytmu Collatza "na trójkach". Rozważmy następujący program. Zakładamy, że program IC startuje z wartościami x, y, z spełniającymi warunek $n \cdot 3^x + y = 2^z \wedge \neg Err$. Jest to warunek *wstępny* obliczeń.

$$IC : \left\{ \begin{array}{l} \text{while } 3^x + y \neq 2^z (* \text{ tj. } n \neq 1 *) \text{ do} \\ \quad \text{if } (\text{odd}(y) \wedge ((x = 0) \text{ or } (y < 3^{x-1}))) \text{ then } Err := \text{true}; \text{ exit fi;} \\ \quad \text{if odd}(y) \text{ then } x := x - 1; \ y := y - 3^x; (*n := 3 * n + 1 *) \\ \quad \text{else } z := z - 1; \ y := y \text{ div } 2; (*n := } \frac{n}{2} *) \text{ fi} \\ \text{od} \end{array} \right\}$$

Obliczenie w strukturze trójek jest zawsze skończone. Wynika to z następującego twierdzenia algorytmicznej teorii liczb naturalnych \mathcal{ATN} .

$$\mathcal{ATN} \vdash \forall x \{\text{while } x \neq 0 \text{ do } x := x - 1 \text{ od}\}(x = 0)$$

Co prawda, może się zdarzyć, że obliczenie algorytmu *IC* zakończy się niepowodzeniem, $Err = \text{true}$.

4.4. Własności programu IC

Fakt 1. *Każde obliczenie programu IC jest skończone i albo jest sukces i osiągnięto jedynkę albo jest błąd i obliczenia algorytmu IC nie można kontynuować.*

$$\mathfrak{N} \models \forall n (n \cdot 3^x + y = 2^z) \implies \{IC\} \left(\underbrace{(3^x + y = 2^z)}_{\langle x, y, z \rangle \asymp 1} \vee \underbrace{(\text{odd}(y) \wedge (x = 0 \vee y < 3^{x-1}))}_{Error} \right)$$

Błąd o jakim tu mowa to niemożność kontynuowania obliczeń. Jest tak gdy y jest nieparzyste i $y < 3^{x-1}$ lub gdy y jest nieparzyste i $x = 0$. Pierwszy błąd łatwo wyeliminować, zauważ, że trójką $\langle x, y, z \rangle$ reprezentuje tę samą liczbę co trójką $\langle x, y+2^z, z+1 \rangle$. Wystarczy więc w sytuacji gdy y jest nieparzyste i $y < 3^{x-1}$ zastąpić trójkę $\langle x, y, z \rangle$ przez trójkę $\langle x, y+2^z, z+1 \rangle$. Zniknie też ryzyko wystąpienia błędu drugiego rodzaju. Upoważnia to nas do sformułowania następującej uwagi.

Uwaga 2. Jeśli obliczenie algorytmu IC rozpoczynające się od pewnej trójki $\langle x, y, z \rangle$ kończy się po k iteracjach błędem $Err = true$ to obliczenie algorytmu IC rozpoczynającego się od trójki $\langle x+1, 3y+2^z, z+2 \rangle$ będzie dłuższe o co najmniej dwie iteracje. Obie te trójki reprezentują tę sama liczbę n .

4.5. Program IIC

Powyzsze uwagi prowadzą do napisania programu poszukujacego dla liczby n dobrej trójki ją reprezentujacej.

```

read(n);
Let  $z = (\mu r)(2^r \geq n)$ .
 $x, x_s := 0; z_s := z; y, y_s := 2^z - n;$ 
Err := false;
while  $3^{x_s} + y_s \neq 2^{z_s}$  do
    while  $3^x + y \neq 2^z$  do
        if odd(y)  $\wedge (x = 0 \vee y < 3^{x-1})$ 
            then Err:=true; exit fi;
        if odd(y) then  $y := y - 3^{x-1}; x := x - 1$ 
            else  $y := y/2; z := z - 1$  fi;
    od;
    if Err then
         $x, x_s := x_s + 1; z, z_s := z_s + 2; y, y_s := 2^{z_s} + 3 \cdot y_s;$ 
        Err := false;
    else exit fi;
od

```

IIC:

Nietrudno zauważyć, że program IIC pomija trójki prowadzące do błędu".

Remark 4.1. Dla danej liczby n program IIC ma obliczenie nieskończone wtedy i tylko wtedy gdy program Cl ma obliczenie nieskończone.

Program *IIC* dokonuje systematycznego przeszukiwania w zbiorze T_n i ewentualnie zwraca najmniejszą trójkę "dobra", tj. taką, że obliczenie programu *IC* jest udane.

5. Proof of Collatz theorem

Our plan may be summarized in four points:

- (i) We proved that the sentence $\forall_n \exists_{x,y,z} n \cdot 3^x + y = 2^z$ is a theorem of the elementary theory \mathcal{T}^+ of natural numbers with addition (Presburger theory). See Appendix B, page 28. Hence, this sentence holds true in any model of the theory.
- (ii) In Appendix C, page 29, we show infinite computations of Collatz's algorithm in a non-standard (non-Archimedean) model of Presburger arithmetic *Ar*. The model is computable and programmable. This example is not a complete counter-example against Collatz conjecture. It only shows that the Collatz conjecture is not a theorem of elementary theory of natural numbers with addition or any other elementary theory.
- (iii) We proved that there is a model \mathfrak{M} of the \mathcal{T}^+ theory such, that it contains an element ε , for which the Collatz algorithm has an infinite computation, c.f. lemma 5.1. *We do not assume* that this model contains unreachable elements.
- (iv) We show that in any model of \mathcal{T}^+ theory, if for a certain n element, the computation of the Collatz algorithm is infinite, then the model is not isomorphic to the standard model of natural numbers (for it contains unreachable elements).

From this we conclude, that if a model has no unreachable elements, then there are no infinite computations.

5.1. Structure in which some element has an infinite Collatz computation.

We have known that in the non-standard model of Presburger arithmetic \mathfrak{M} unreachable elements have infinite Collatz computations. The model is described in the literature, see [Grz71]. We provide examples of infinite Collatz computations and a definition

of this structure in a programming language in section 9, Appendix C. Now, we are going to show that there are non-Collatz elements without assuming that they are unreachable elements.

We construct an elementary theory \mathcal{T}^+ which is an extension of the theory of \mathcal{T} (i.e. Presburger arithmetic) in a way that permits to show an element different from any number that occurs in Collatz tree.

The extension of theory \mathcal{T} is made in three steps

1. (*language*) we add a new constant ε to the alphabet and correspondingly we extend the sets of term and of formulas of the language of theory \mathcal{T} .
2. (*logic*) the operation of consequence remains the same, remember the sets of terms and of formulas are bigger,
3. (*axioms of data structure*) to the set Ax of Presburger's axioms we add an infinite set of sentences Z .

Our intention is to prove the following fact. An assumption that for some element ε the Collatz computation is infinite, does not lead to a contradiction with axioms of elementary theory of addition of natural numbers (i.e. the Presburger's theory).

As a natural reflex, we would like to add the negation of the instance of halting formula for $n = \varepsilon$ to the axioms of of \mathcal{T} theory. However the formula

$$\neg\{n \leftarrow \varepsilon\} \left\{ \begin{array}{l} \text{while } n \neq 1 \text{ do} \\ \quad \text{if } even(n) \\ \quad \text{then } n \leftarrow n \text{ div } 2 \\ \quad \text{else } n \leftarrow 3n + 1 \\ \quad \text{fi} \\ \text{od} \end{array} \right\} (n = 1)$$

does not belong to the language of elementary theory of \mathcal{T} . Moreover, the following formula 18 that expresses the same looping property of computation of Collatz algorithm does not belong to

the language of first-order theory *Ar.*

$$\{n \leftarrow \varepsilon\} \cap \left\{ \begin{array}{l} \text{if } n \neq 1 \text{ then} \\ \quad \text{if } n \bmod 2 = 0 \\ \quad \text{then } n \leftarrow n \text{ div } 2 \\ \quad \text{else } n \leftarrow 3n + 1 \\ \quad \text{fi} \\ \text{fi} \end{array} \right\} (n \neq 1) \quad (18)$$

However, for every algebraic structure \mathfrak{A} the above formula (18) is valid in \mathfrak{M} if and only if every formula of the following scheme is valid in \mathfrak{A}

$$\begin{aligned} & \{n \leftarrow \varepsilon\}(n \neq 1), \\ & \{n \leftarrow \varepsilon\} \left\{ \begin{array}{l} \text{if } n \neq 1 \text{ then} \\ \quad \text{if } \text{even}(n) \text{ then } n \leftarrow n \text{ div } 2 \\ \quad \text{else } n \leftarrow 3n + 1 \text{ fi} \\ \text{fi} \end{array} \right\} (n \neq 1), \\ & \dots \\ & \{n \leftarrow \varepsilon\} \left\{ \begin{array}{l} \text{if } n \neq 1 \text{ then} \\ \quad \text{if } \text{even}(n) \text{ then } n \leftarrow n \text{ div } 2 \\ \quad \text{else } n \leftarrow 3n + 1 \text{ fi} \\ \text{fi} \end{array} \right\}^i (n \neq 1), \\ & \dots \end{aligned}$$

Each of these formulas is equivalent to certain first-order formula ϑ , such that the formula does not contain any algorithm. One can verify this claim making use of axioms of assignment instruction e.g. $\{n \leftarrow \varepsilon\}(n > 1) \equiv (\varepsilon > 1)\}$, conditional instruction and composition of programs. We illustrate our claim by the following equivalence.

$$\begin{aligned} & \{n \leftarrow \varepsilon\} \{\text{if } P(n) \text{ then } n \leftarrow n \text{ div } 2 \text{ else } n \leftarrow 3n + 1 \text{ fi}\}(n \neq 1) \equiv \\ & \quad \left((P(\varepsilon) \wedge \varepsilon \neq 2) \vee \underbrace{(\neg P(\varepsilon) \wedge 3\varepsilon + 1 \neq 1)}_{\text{false}} \right) \end{aligned}$$

Continuing, we obtain the following formulas

Formuła
$(o(\varepsilon) \wedge \varepsilon \neq 1)$
$(e(\varepsilon) \wedge o(\frac{\varepsilon}{2}) \wedge \varepsilon \neq 2)$
$(e(\varepsilon) \wedge e(\frac{\varepsilon}{2}) \wedge o(\frac{\varepsilon}{4}) \wedge \varepsilon \neq 4)$
$(e(\varepsilon) \wedge e(\frac{\varepsilon}{2}) \wedge e(\frac{\varepsilon}{4}) \wedge o(\frac{\varepsilon}{8}) \wedge \varepsilon \neq 8)$
$(e(\varepsilon) \wedge e(\frac{\varepsilon}{2}) \wedge e(\frac{\varepsilon}{4}) \wedge e(\frac{\varepsilon}{8}) \wedge o(\frac{\varepsilon}{16}) \wedge \varepsilon \neq 16)$
$\left(\begin{array}{l} e(\varepsilon) \wedge e(\frac{\varepsilon}{2}) \wedge e(\frac{\varepsilon}{4}) \wedge e(\frac{\varepsilon}{8}) \wedge e(\frac{\varepsilon}{16}) \wedge o(\frac{\varepsilon}{32}) \wedge \varepsilon \neq 32 \vee \\ o(\varepsilon) \wedge e(3\varepsilon + 1) \wedge e(\frac{3\varepsilon + 1}{2}) \wedge e(\frac{3\varepsilon + 1}{4}) \wedge e(\frac{3\varepsilon + 1}{8}) \wedge o(\frac{3\varepsilon + 1}{16}) \wedge \varepsilon \neq 5 \end{array} \right)$
$\left(\begin{array}{l} (e(\varepsilon) \wedge e(\frac{\varepsilon}{2}) \wedge e(\frac{\varepsilon}{4}) \wedge e(\frac{\varepsilon}{8}) \wedge e(\frac{\varepsilon}{16}) \wedge e(\frac{\varepsilon}{32}) \wedge o(\frac{\varepsilon}{64}) \wedge \varepsilon \neq 64 \vee \\ e(\varepsilon) \wedge o(\frac{\varepsilon}{2}) \wedge e(\frac{3\varepsilon + 1}{2}) \cdots \wedge \varepsilon \neq 10 \end{array} \right)$
$\left(\begin{array}{l} e(\varepsilon) \wedge e(\frac{\varepsilon}{2}) \wedge e(\frac{\varepsilon}{4}) \wedge e(\frac{\varepsilon}{8}) \wedge e(\frac{\varepsilon}{16}) \wedge e(\frac{\varepsilon}{32}) \wedge e(\frac{\varepsilon}{64}) \wedge o(\frac{\varepsilon}{128}) \wedge \varepsilon \neq 128 \vee \\ o(\varepsilon) \wedge e(3\varepsilon + 1) \wedge e(\frac{3\varepsilon + 1}{2}) \cdots \wedge \varepsilon \neq 21 \vee \\ e(\varepsilon) \wedge o(\frac{\varepsilon}{2}) \wedge e(\frac{3\varepsilon + 1}{2}) \cdots \wedge \varepsilon \neq 20 \vee \\ o(\varepsilon) \wedge e(3\varepsilon + 1) \wedge o(\frac{3\varepsilon + 1}{2}) \cdots \wedge \varepsilon \neq 3 \end{array} \right)$
...

Horizontal lines separate formulas corresponding to different levels of Collatz tree. We define the set Z as containing all formulas $\varepsilon \neq k$ such,,that the expression $n = k$ occurs in halting formula of Collatz algorithm, k is number. Hence, the set Z contains the sentences $\{\varepsilon \neq 1, \varepsilon \neq 2, \varepsilon \neq 4, \varepsilon \neq 8, \varepsilon \neq 16, \varepsilon \neq 32, \varepsilon \neq 5, \varepsilon \neq 64, \varepsilon \neq 10, \varepsilon \neq 128, \varepsilon \neq 20, \varepsilon \neq 21, \varepsilon \neq 3, \dots\}$

Na zbiór Ax' aksjomatów nowej teorii składają się: aksjomaty Ax teorii Presburgera (por. Dodatek A, strona 26) oraz nieskończony zbiór Z .

$$Ax' = Ax \cup Z$$

Udowodnimy, że zbiór Ax' jest niesprzeczny. Zaczniemy od wykazania, że każdy skończony podzbiór Ax_0 zbioru Ax' jest niesprzeczny. Wykażemy mianowicie, że zbiór Ax_0 posiada model w standardowej strukturze \mathfrak{N} liczb naturalnych z dodawaniem. Nasze zadanie ogranicza się do podania właściwej interpretacji stałej: ε . Niech Z_0 będzie dowolnym skończonym podzbiorem zbioru Z . Zbiór zdań $Ax \cup Z_0$ jest niesprzeczny. Aby to wykazać weźmy jako

wartość stałej ε jakąś liczbę l większą od każdej liczby k występującą w zbiorze zdań Z_0 . Taki wybór zapewnia prawdziwość każdego zdania ze zbioru Z_0 .

We proved that every finite subset of the set Ax' is consistent. By the compactness theorem on first-order logic⁴, we obtain

Lemma 5.1. The set Ax' is a consistent set of formulas.

Now, we apply the model existence theorem⁵, which reads: *for every consistent set S of first-order formulas there exists an algebraic structure \mathfrak{A} such that it is a model of the set S , i.e. for every formula $\sigma \in S$ the formula is valid in the structure \mathfrak{A} .*

In this way we proved the following lemma.

Lemma 5.2. There is an algebraic structure \mathfrak{M} , such that every sentence of the set Ax' is valid in it.

Corollary 5.1. The execution of Collatz algorithm in structure \mathfrak{M} that starts with value of variable n equal ε , $v(n) = \varepsilon$ is infinite.

For the structure \mathfrak{M} is a model of the set Z .

Corollary 5.2. Element ε of structure \mathfrak{M} does not belong to the Collatz tree DC .

It remains to be proved that every standard natural number has a finite Collatz computation.

This it is equivalent to the following statement *for every element n if its Collatz computation is infinite than the element n is non-standard (i.e. unreachable) element of a model of elementary theory of natural numbers.*

⁴Compactness theorem *If every finite subset of a set S of formulas is consistent, then the set S is consistent too.*

⁵of first-order logic

5.2. Infinite Collatz computation require unreachable elements

In our proof we shall imitate the proof of lemma 5.1 . Let ε be an arbitrary element such, that the execution of Collatz algorithm is infinite. (i.e. element ε is non-Collatz one). It is evident that the execution of algorithm *IIC* (c.f. 4.5) for element ε is infinite, too. For any triple $\langle x, y, z \rangle$ of standard, natural numbers the execution of program *IC* terminates with the error *Err*.

Yet, the execution of Collatz algorithm for the element ε is infinite. Our nearest goal is to prove that there exist three elements c_x, c_y, c_z such that 1° the equality $\varepsilon \cdot 3^{c_x} + c_y = 2^{c_z}$ holds and 2° the computation of algorithm *IC* is infinite.

We extend the language of the elementary theory \mathcal{T}^+ badding to it four constants: c_n, c_x, c_y, c_z . We fix the value of constant c_n assuming that the equality $c_n = \varepsilon$ is an axiom of new theory. We are going to show that the following set of first-order sentences $Ax' = Ax \cup Df \cup U \cup Y$ is consistent. Sets Ax and Z were described earlier. Set U contains one formula $c_n \cdot 3^{c_x} + c_y = 2^{c_z}$. Set Df contains definitions of useful operations and relations $P2, P3, 3x, \text{div2}, \text{even}, \text{odd}$, c.f. section 7.

Set Y contains all first-order sentences quivalent to the algorithmic sentences of the form

$$\left\{ \begin{array}{l} x := c_x; \\ y := c_y; \\ z := c_z; \\ Err := \text{false} \end{array} \right\} \left\{ \begin{array}{l} \text{if } 3^x + y \neq 2^z \text{ then} \\ \quad \text{if } (\text{odd}(y) \wedge (x = 0 \vee y < 3^{x-1})) \text{ then } Err := \text{true} \\ \quad \text{else} \\ \quad \quad \text{if } \text{odd}(y) \text{ then} \\ \quad \quad \quad x := x - 1; y := y - 3^x \\ \quad \quad \quad \text{else} \\ \quad \quad \quad y := y \div 2; z := z - 1 \\ \quad \quad \quad \text{fi} \\ \quad \quad \text{fi} \\ \quad \text{fi} \end{array} \right\}^i (\neg Err)$$

where $i = 0, 1, 2, \dots$

In other words Y is a sequence $Y = \{y_0, y_1, y_2, \dots\}$.

Sentence y_i expresses the following semantical property: *Let s be an execution of algorithm IC that starts with triple $\langle c_x, c_y, c_z \rangle$. If during this execution, after i steps and reached the state x, y, z then if the current state does not represents the number 1, then in the next step no error Err will be raised.*

In other words the sentence y_i excludes the risk of error Err in $i+1$ -th step of execution.

As we did it earlier, we show that every finite subset $Ax_0 \subset Ax'$ is consistent. We shall use observations made in section 4.5, that concern the algorithm *IIC*. Let Y_0 be any finite subset of the set Y of sentences. Let i be the largest number of iterations mentioned in the set . From the properties of algorithm *IIC*we know that there exists triple $\langle x, y, z \rangle$ of natural numbers such, that during execution of i iterations of program \bar{K} no error *Err* will occur.

Hence, we can apply the compactness theorem and assert that the whole set Ax' is consistent.

By completeness theorem we infer that there is a triple $\langle x, y, z \rangle$ such ,that for every natural number i after execution of i iterations of the while instruction in algorithm *IC* the next iteration can be executed without risk of *Err* error. I.e. the condition θ mentioned in the lemma 4.3 is satisfied. Let us cite it here

$$\theta : \left(\begin{array}{l} (\varepsilon \cdot 3^{cx} + cy = 2^{cz}) \wedge \\ \left\{ \begin{array}{l} x := cx; \\ y := cy; \\ z := cz \end{array} \right\} \cap \left\{ \begin{array}{l} \text{if } 3^x + y \neq 2^z \\ \text{then} \\ \quad \text{if odd}(y) \\ \quad \text{then } x, y := x - 1, y - 3^{x-1} \\ \quad \text{else } z, y := z - 1, y \div 2 \\ \text{fi} \\ \text{fi} \end{array} \right\} \left(\begin{array}{l} \text{odd}(y) \implies (x > 0 \wedge y > 3^{x-1}) \end{array} \right) \end{array} \right)$$

Hence, two formulas hold: the formula θ and $\neg\{n := \varepsilon; Cl\}(n = 1)$. From this conjunction we deduce

Corollary 5.3. If the condition θ holds and the execution of program *Cl* for $n = \varepsilon$ is infinite, then the computation of program *IC* starting with the triple $\langle c_x, c_y, c_z \rangle$ is infinite too .

Proof:

This is an immediate consequence of the lemma 4.3. □

5.3. Collatz theorem

Let's summarize what we know:

- If the Collatz computation for an element n is finite, then there is a triple $\langle x, y, z \rangle$ such that $n \cdot 3^x + y = 2^z$ and the computation of algorithm *IC* for this triple is finite.
- If for some triple $\langle x, y, z \rangle$, the following equality holds $n \cdot 3^x + y = 2^z$ and the computation of algorithm*IC* is finite, then the computation of the Collatz algorithm for n is finite.

- if in a model \mathfrak{M} of the elementary theory of natural numbers with addition (i.e. Presburger theory) an element n is unreachable, then the computation of the Collatz algorithm for n is infinite.
- There is a structure \mathfrak{M} , model of Presburger arithmetic and an element ε such that, the computation of the Cl algorithm is infinite.
- Let an algebraic structure \mathfrak{A} be a model of Presburger arithmetic. Let n be any element for which the computation of the Collatz algorithm is infinite. There are three elements: $\langle x, y, z \rangle$ such, that the equality $n \cdot 3^x + y = 2^z$ holds and the computation of the IC algorithm for this triple is infinite.

From these facts we derive the following proposition.

Theorem 5.1. (Collatz 1937)

For every natural number n , the execution of Collatz algorithm is finite.

Proof:

It follows from the facts enumerated above that if for some element n of the \mathfrak{A} structure, which is a model of the elementary theory of the addition of natural numbers, the calculation of the Collatz algorithm is infinite, then the structure is a non-standard model of the theory. Hence, by transposition, we obtain Collatz theorem.

□

6. Podsumowanie

Nietrudno zauważyć, że przedstawiony dowód jest okrężny. Nie potrafimy, na razie, przedstawić dowodu wyprowadzonego wprost w algorytmicznej teorii $AT\mathcal{N}$ z aksjomatów tej teorii lub np. z prawa Archimedesa, które jest twierdzeniem teorii $AT\mathcal{N}$.

Kolejne zadanie to oszacowanie kosztu algorytmu Collatza, wiemy, że obliczenia są skończone, ale nie potrafimy oszacować ich długosci. Natomiast koszt odpowiedzi na pytanie czy liczba n jest elementem drzewa Collatza jest stały $O(1)$. Z twierdzenia Collatza potrafimy jednak wyprowadzić kilka ciekawych wniosków, o czym napiszemy osobno.

Podziękowania

Andrzej Szałas znalazł błąd w innej, wcześniejszej pracy na temat algorytmu Collatza. Wiktor Dańko zwrócił naszą uwagę na lukę w analizie obliczeń trójkowych. Ludwik Czaja i Marek Warpechowski przekazali nam swe wątpliwości, dzięki nim powstała obecna wersja pracy.

Wszystkie usterki i błędy są naszego autorstwa.

7. Dodatek A – Kilka faktów o elementarnej teorii dodawania

W rozdziale 4 zaobserwowałyśmy parę pozytycznych faktów o trójkach reprezentujących liczby naturalne.

Rozważać będziemy następującą teorię T_+ , por. [Grz71] str. 239 i następne.

Definicja 1. Teoria $T_+ = \langle \mathcal{L}, \mathcal{C}, Ax \rangle$ jest układem trzech przedmiotów:

\mathcal{L} jest językiem pierwszego rzędu. Na alfabet tego języka składają się: zbiór V zmiennych, znaki operacji: $0, S, +$, znak relacji równości $=$, znaki funkторów logicznych i kwantrifikatorów, znaki pomocnicze, m. in nawiasy..

Zbiór wyrażeń poprawnie zbudowanych to suma teoriomnoścowa zbioru termów T i zbioru formuł F .

Zbiór termów T jest to najmniejszy zbiór napisów zawierający zbiór zmiennych V i napis 0 i zamknięty ze względu na reguły: jeśli dwa napisy τ_1 oraz τ_2 są termami to termem jest też napis postaci $(\tau_1 + \tau_2)$, jeśli napis τ jest termem to napis $S(\tau)$ jest także termem.

Zbiór formuł jest najmniejszym zbiorem napisów zawierającym równości tj. napisy postaci $(\tau_1 = \tau_2)$ i zamkniętym ze względu na reguły: jeśli napisy α oraz β są formułami to formułami są też napisy postaci

$$(\alpha \vee \beta), (\alpha \wedge \beta), (\alpha \implies \beta), \neg \alpha$$

formułami są też napisy postaci

$$\forall_x \alpha, \exists_x \alpha$$

gdzie x jest zmienną, a α jest formuła.

\mathcal{C} jest operacją konsekwencji zdeterminowaną przez przyjęcie aksjomatów logiki pierwszego rzędu (rachunku predykatów) i reguł wnioskowania logiki pierwszego rzędu

Ax jest zbiorem formuł wyliczonych poniżej.

$$\begin{aligned}
 \forall_x x + 1 \neq 0 & \tag{a} \\
 \forall_x \forall_y x + 1 = y + 1 \implies x = y & \tag{b} \\
 \forall_x x + 0 = x & \tag{c} \\
 \forall_{x,y} (y + 1) + x = (y + x) + 1 & \tag{d} \\
 \{\Phi(0) \wedge \forall_x [\Phi(x) \implies \Phi(x + 1)] \implies \forall_x \Phi(x) & \tag{I} \\
 \end{aligned}
 \tag{19}$$

Tu napis $\Phi(x)$ należy zastąpić jakikolwiek formułą. Ostatni wiersz jest więc schematem indukcji.

Do tego zbioru dodajemy aksjomaty definiujące dodatkowe pojęcia

$$\begin{aligned}
 Parz(x) \stackrel{df}{=} \exists_y x = y + y & \tag{p} \\
 x \text{ div } 2 = y \equiv (x = y + y \vee x = y + y + 1) & \tag{D2} \\
 3x \stackrel{df}{=} x + x + x & \tag{3x}
 \end{aligned}$$

W teorii T_+ przeprowadzimy dowody paru faktów znanych z rozdziału 4. Dzięki temu upewnimy się, że fakty te prawdziwe są także w każdym modelu teorii. Natomiast w rozdziale 5 posłużymy się teorią Presburgera

Definicja 2. Teoria $Ar = \langle \mathcal{L}, \mathcal{C}, Ax \rangle$ jest układem trzech przedmiotów:

\mathcal{L} jest językiem pierwszego rzędu. Na alfabet tego języka składają się: zbiór V zmiennych, znaki operacji: $0, +$, znak relacji równości $=$.

Zbiór wyrażeń poprawnie zbudowanych to unia zbioru termów T i zbioru formuł F . Zbiór termów T jest to najmniejszy zbiór napisów zawierający zbiór zmiennych V i napis 0 i zamknięty ze względu na reguły: jeśli dwa napisy τ_1 oraz τ_2 są termami to termem jest też napis postaci $(\tau_1 + \tau_2)$, jeśli napis τ jest termem to napis $S(\tau)$ jest także termem.

\mathcal{C} jest operacją konsekwencji zdeterminowaną przez przyjęcie aksjomatów rachunku predykatów i reguł wnioskowania logiki pierwszego rzędu

Ax jest zbiorem formuł wyliczonych poniżej.

$$\forall_x x + 1 \neq 0 \quad (\text{A})$$

$$\forall_x x \neq 0 \implies \exists_y x = y + 1 \quad (\text{B})$$

$$\forall_{x,y} x + y = y + x \quad (\text{C})$$

$$\forall_{x,y,z} x + (y + z) = (x + y) + z \quad (\text{D})$$

$$\forall_{x,y,z} x + z = y + z \implies x = y \quad (\text{E})$$

$$\forall_x x + 0 = x \quad (\text{F})$$

$$\forall_{x,z} \exists_y (x = y + z \vee z = y + x) \quad (\text{G})$$

$$\forall_x \exists_y (x = y + y \vee x = y + y + 1) \quad (\text{H2})$$

$$\forall_x \exists_y (x = y + y + y \vee x = y + y + y + 1 \vee x = y + y + y + 1 + 1) \quad (\text{H3})$$

.....

$$\forall_x \exists_y \left(\begin{array}{l} x = \underbrace{y + y + \cdots + y}_k \vee \\ x = \underbrace{y + y + \cdots + y}_{k-1} + 1 \vee \\ x = \underbrace{y + y + \cdots + y}_k + \underbrace{1 + 1}_2 \vee \\ \dots \\ x = \underbrace{y + y + \cdots + y}_k + \underbrace{1 + 1 + \cdots + 1}_{k-2} \vee \\ x = \underbrace{y + y + \cdots + y}_k + \underbrace{1 + 1 + \cdots + 1}_{k-1} \end{array} \right) \quad (\text{Hk})$$

....

Przypomnijmy parę faktów

F1. Teoria T_+ jest elementarnie równoważna teorii Ar . [Pre29, Sta84]

F2. Teoria Ar jest rozstrzygalna. [Pre29].

F3. Złożoność teorii Ar , czyli koszt udowodnienia, że dane zdanie α

jest twierdzeniem (lub jego negacją) jest rzędu podwójnie wykładniczego $O(2^{2^n})$. [FR79].

F4. Teorie T_+ oraz Ar mają modele niestandardowe, tj. modele zawierające elementy nieosiągalne, zob Dodatek C, strona 29.

8. Dodatek B– dowód twierdzenia 4.1

W tym rozdziale wykażemy, że zdanie *dla każdego n istnieją x, y, z takie, że $n \cdot 3^x + y - 2^z$* jest twierdzeniem teorii T_+ dodawania. Elemen-tarna teoria T_+ dodawania liczb naturalnych została przypomniana w dodatku A.

Działania mnożenia i potęgowania są niedostępne w teorii T_+ . Ale nie są niezbędne do osiągnięcia naszego celu.

Teorię T_+ wzbogacamy o dwie funkcje $P2(\cdot)$ oraz $P3(\cdot\cdot)$. zdefiniowane w ten sposób

$$\begin{array}{ll} P2(0) = 1 & P3(y, 0) = y \\ P2(x+1) = P2(x) + P2(x) & \left| \begin{array}{l} P3(y, x+1) = P3(y, x) + P3(y, x) + P3(y, x) \end{array} \right. \end{array}$$

Lemat 3. Powyższe definicje są poprawne, tzn. twierdzeniami teorii T_+ wzbogaconej w ten sposób są zdania $\forall_x \exists_y P2(x) = y$ i $\forall_{x,y,z} P2(x) = y \wedge P2(x) = z \implies y = z$.

$$T_+ \vdash \forall_x \exists_y P2(x) = y \quad \mathbf{i}$$

$$T_+ \vdash \forall_{x,y,z} P2(x) = y \wedge P2(x) = z \implies y = z.$$

Podobnie twierdzeniami wzbogaconej teorii T_+ są zdania $\forall_{y,x} \exists_z P3(y, x) = z$ i $\forall_{y,x,z,u} P3(y, x) = z \wedge P3(y, x) = u \implies z = u$.

Dowód przebiega przez indukcję względem zmiennej x .

Potrzebna nam będzie następująca definicja relacji mniejszości $a < b \stackrel{df}{=} \exists_c a + S(c) = b$. Wykorzystując definicje funkcji $P2$ oraz $P3$ napiszemy wyrażenie $P3(n, x) + y = P2(z)$.

Lemma 8.1. Następujące zdanie jest twierdzeniem wzbogaconej teorii T_+

$$\forall_n \exists_{x,y,z} P3(n, x) + y = P2(z)$$

Najpierw przez indukcję dowodzimy, że $T_+ \vdash \forall_n n < 2^n$. A dokładniej, $T_+ \vdash \forall_n n < P2(n)$. Łatwo sprawdzić, że $T_+ \vdash 0 < P2(0)$. Założymy, że $T_+ \vdash \forall_n \{n < P2(n)\}$. Nierówność $n + 1 < P2(n + 1)$ wynika z dwu

nierówności $T_+ \vdash n < P2(n)$ i $\mathcal{T} \vdash 1 < P2(n)$.

W podobny sposób uzyskamy $T_+ \vdash P3(n, x) < P2(z) \wedge (z = n + x + x)$
 Wynika stąd, że $T_+ \vdash \forall_n \exists_{x,y,z} P3(n, x) + y = P2(z)$.
 Właściwie wykazaliśmy, że $\forall_{n,x} \exists_{y,z} P3(n, x) + y = P2(z)$

Lemma 8.2. Niech \mathfrak{M} będzie jakimkolwiek modelem arytmetyki Presburgera. Element n jest osiągalny wtedy i tylko wtedy gdy istnieje taka trójka reprezentująca element n , tj. taka, że zachodzi równość $P3(n, x) + y = P2(z)$ czyli $n \cdot 3^x + y = 2^z$ i liczby x, y, z są osiągalne.

Proof:

Jeśli spełnione są formuły

$$\{q := 0; \text{while } q \neq x \text{ do } q := q + 1 \text{ od}\}(x = q),$$

$$\{q := 0; \text{while } q \neq y \text{ do } q := q + 1 \text{ od}\}(y = q),$$

$$\{q := 0; \text{while } q \neq z \text{ do } q := q + 1 \text{ od}\}(z = q)$$

i ponadto zachodzi równość

$P3(n, x) + y = P2(z)$ to łatwo sprawdzić, że spełniona jest formuła $\{t := 0; \text{while } n \neq t \text{ do } t := t + 1 \text{ od}\}(t = n)$ □

9. Appendix C - an example of an infinite computation

At this point, we'll remind you of a few facts that are less known to the IT community. In Appendix A we described the Ar theory of addition of natural numbers. The only functor in the language of this theory is $+$, we also have two constants 0 and 1 and the predicate of equality $=$. Now, we will program the algebraic structure \mathfrak{M} , which is a model of this theory, i.e. all axioms of theory Ar are true in the structure \mathfrak{M} . First we will describe this structure as mathematicians do, then we will write a class (ie a program module) implementing this structure. medskip

9.1. Mathematical description of the structure

\mathfrak{M} is an algebraic structure

$$\mathfrak{M} = \langle M; \mathbf{0}, \mathbf{1}, \oplus; = \rangle \quad (\text{NonStandard})$$

such that M is a set of pairs $\langle k, w \rangle$ where element $k \in Z$ is an integer, element w is a rational, non-negative number and the following requirements are lsatisfied:

- (i) for each element $\langle k, w \rangle$ if $w = 0$ then $k \geq 0$,
- (ii) the meaning of the constant 0 is $\langle 0.0 \rangle$,
- (iii) the meaning of constant 1 is $\langle 1.0 \rangle$,
- (iv) the operation \oplus of addition is determined as follows

$$\langle k, w \rangle \oplus \langle k', w' \rangle \stackrel{df}{=} \langle k + k', w + w' \rangle.$$

Lemma 9.1. The algebraic structure \mathfrak{M} is a model of *Ar* theory.

The reader will check that each axiom of the *Ar* theory is a sentence true in the structure \mathfrak{M} .

The structure \mathfrak{M} is not a model of the *ATN*, algorithmic theory of natural numbers. Elements of the structure $\langle k, w \rangle$, such as $w \neq 0$ are *unreachable*. i.e. for each element $x_0 = \langle k, w \rangle$ such that $w \neq 0$ the following condition holds

$$\neg\{y := 0; \text{while } y \neq x_0 \text{ do } y := y + 1 \text{ od}\}(y = x_0)$$

The subset $\mathfrak{N} \subset \mathfrak{M}$ composed of only those elements for which $w = 0$ is a model of the theory *ATN*. The elements of the structure \mathfrak{N} are called *reachable*. A very important theorem of the foundations of mathematics is

Fact 9.1. The structures \mathfrak{N} and \mathfrak{M} are not isomorphic. See [Grz71], p. 256.

As we will see in a moment, this fact is also important for IT specialists.

9.2. Definition in programming language

Perhaps you have already noticed that the \mathfrak{M} is computable. The following is a class that implements the structure \mathfrak{M} . The implementation uses the integer type, we do not introduce rationalNumbers explicitly.

```

unit StrukturaM: class;
  unit Elm: class(k,li,mia: integer);
    begin
      if mia=0 then raise Error fi;
      if li * mia <0 then raise Error fi;
      if li=0 and k<0 then raise Error fi;
    end Elm;
    add: function(x,y:Elm): Elm;
    begin
      result := new Elm(x.k+y.k, x.li*y.mia+x.mia*y.li, x.mia*y.mia )
    end add;
    unit one : function:Elm; begin result:= new Elm(1,0,2) end one;
    unit zero : function:Elm; begin result:= new Elm(0,0,2) end zero;
    unit eq: function(x,y:Elm): Boolean;
    begin
      result := (x.k=y.k) and (x.li*y.mia=x.mia*y.li )
    end eq;
  end StrukturaM

```

The following lemma expresses the correctness of the implementation

Lemma 9.2. The set of Elm objects with the *add* operation is a model of the *Ar* theory

9.3. Infinite Collatz algorithm computation

How to execute the Collatz algorithm in StructuraM? It's easy.

```

pref StrukturaM block
  var n: Elm;
  unit odd: function(x:Elm): Boolean; ... result:=(x.k mod 2)=1 ... end odd;
  unit div2: function(x:elm): Elm; ...
begin
  n:= new Elm(8,1,2);
  while not eq(n,one) do
    if odd(n) then
      n:=add(n,add(n,add(n,one))) else n:= div2(n)
    fi
  od
end block;

```

Below we present the computation of Collatz algorithm for $n = \langle 8, \frac{1}{2} \rangle$.

$$\langle 8, \frac{1}{2} \rangle, \langle 4, \frac{1}{4} \rangle, \langle 2, \frac{1}{8} \rangle, \langle 1, \frac{1}{16} \rangle, \langle 4, \frac{3}{16} \rangle, \langle 2, \frac{3}{32} \rangle, \langle 1, \frac{3}{64} \rangle, \langle 4, \frac{9}{64} \rangle, \langle 2, \frac{9}{128} \rangle, \dots$$

None of the elements of the above sequence is a standard natural number. Each of them is unreachable. It is worth looking at an example of another calculation. Will something change when we assign n a different object? e.g. $n := \text{new Elm}(19, 2, 10)$?

$$\begin{aligned} &\langle 19, \frac{10}{2} \rangle, \langle 58, \frac{30}{2} \rangle, \langle 29, \frac{30}{4} \rangle, \langle 88, \frac{90}{4} \rangle, \langle 44, \frac{90}{8} \rangle, \langle 22, \frac{90}{16} \rangle, \langle 11, \frac{90}{32} \rangle, \langle 34, \frac{270}{32} \rangle, \langle 17, \frac{270}{64} \rangle, \\ &\langle 52, \frac{810}{64} \rangle, \langle 26, \frac{405}{64} \rangle, \langle 13, \frac{405}{128} \rangle, \langle 40, \frac{1215}{128} \rangle, \langle 20, \frac{1215}{256} \rangle, \langle 10, \frac{1215}{256} \rangle, \langle 5, \frac{1215}{512} \rangle, \langle 16, \frac{3645}{512} \rangle, \langle 8, \frac{3645}{1024} \rangle, \\ &\langle 4, \frac{3645}{2048} \rangle, \langle 2, \frac{3645}{4096} \rangle, \langle 1, \frac{3645}{8192} \rangle, \langle 4, \frac{3*3645}{8192} \rangle, \langle 2, \frac{3645*3}{2*8192} \rangle, \langle 1, \frac{3*3645}{4*8192} \rangle, \langle 4, \frac{9*3645}{4*8192} \rangle, \dots \end{aligned}$$

And one more computation.

$$\begin{aligned} &\langle 19, 0 \rangle, \langle 58, 0 \rangle, \langle 29, 0 \rangle, \langle 88, 0 \rangle, \langle 44, 0 \rangle, \langle 22, 0 \rangle, \langle 11, 0 \rangle, \langle 34, 0 \rangle, \langle 17, 0 \rangle, \langle 52, 0 \rangle, \langle 26, 0 \rangle, \\ &\langle 13, 0 \rangle, \langle 40, 0 \rangle, \langle 20, 0 \rangle, \langle 10, 0 \rangle, \langle 5, 0 \rangle, \langle 16, 0 \rangle, \langle 8, 0 \rangle, \langle 4, 0 \rangle, \langle 2, 0 \rangle, \langle 1, 0 \rangle \end{aligned}$$

Corollary 9.1. The structure \mathfrak{M} , which we have described in two different ways, is the model of the T_+ theory (you can also say that this structure implements the specification given by the axioms of the Ar theory), with the non-obvious presence of unreachable elements in it.

Another observation

Corollary 9.2. The halting property of the Collatz algorithm cannot be proved from the axioms of the T_+ theory, nor from the Ar theory.

The reader may wish to construct the computation that starts with $\langle 8, \frac{1}{7} \rangle$.

References

- [FR79] Jeanne Ferrante and Charles W. Rackoff. The Computational Complexity of Logical Theories. Springer Verlag, Heidelberg, 1979.
- [Grz71] Andrzej Grzegorczyk. Zarys Arytmetyki Teoretycznej. PWN, Warszawa, 1971, pp.311.

- [MS87] Grażyna Mirkowska and Andrzej Salwicki. Algorithmic Logic. PWN, Warszawa, 1987, pp.372. http://lem12.uksw.edu.pl/wiki/Algorithmic_Logic, 1987. [Online; accessed 7-August-2017].
- [Pre29] Mojżesz Presburger. Über die Vollständigkeit eines gewissen Systems der Aritmetik. Comptes Rendus du 1-er Congrès des Mathematiciens des Pays Slaves, Varsovie, 1929, 95-101.
- [Sta84] Ryan Stansifer. Presburger's Article on Integer Arithmetic: Remarks and Translation. Technical Report TR84-639, Cornell University, 1984.